

# BSM Parton Shower in Herwig7

1. Generalized angular-order parton showers in Herwig 7  
[Lee, Masouminia, Seymour, Yang, JHEP 08 (2024) 064]
2. Searching for New Physics Inside Jets with the Herwig 7  
Generalised Parton Shower  
[Lee, Kim, Masouminia, Seymour, Yang]

Joon-Bin Lee<sup>1</sup> • M.R. Masouminia<sup>3</sup> • Taehee Kim<sup>1</sup> • Un-ki Yang<sup>1</sup> • Michael H. Seymour<sup>2</sup>

<sup>1</sup> Seoul National University

<sup>2</sup> University of Manchester

<sup>3</sup> IPPP, Durham University




MCnet project




Marie Curie Actions






# Event Generator = Translator

Ultra-high dimension  
integration



“MC method”



ME event generators


- MadGraph5, POWHEG, Sherpa, ...

General purpose (Parton Shower) event generators

- Pythia8, Herwig7, ...

# Event Simulation in pp Collisions

@ low  $Q^2$ , running  $\alpha_s$  = infinite number of diagrams



## QCD simulation

- > How to control **soft & collinear physics (divergences)**?
- > For processes with separate scales  $Q \gg Q_0$ , higher-order corrections are enhanced by large logarithms

$$\alpha_s^n \ln^m Q/Q_0$$

- > Hadron physics needs **logarithmic resummation**  
= **Parton Shower**

## Angular-order shower program (Herwig7)

- > Handle colour coherence
- > **QCD shower (1988)** → **QED (2006)** → **EW (2022)** → **“BSM (2024)”**  
extensions in the parton shower regime

[Lee, Masouminia, Seymour, Yang, JHEP 08 (2024) 064]

# Parton Shower Algorithms


Comparing Approaches: Angular-Ordering vs. Dipole Shower

## Herwig7

Angular-Ordered Shower

### Strength

- Color Coherence** Captures the "Chudakov effect" naturally via angular ordering property (simple).



- Shower History** Perfectly tracks shower history, making it ideal for recursive BSM implementations.

- BSM Suitability** Splitting functions are more intuitive. Excellent for implementing new radiation.


VS

## Pythia8

p<sub>T</sub>-Ordered Dipole Shower

### Strength

- Color Coherence** Dipole algorithm (with p<sub>T</sub> ordering).



- Adoption** CMS default parton shower generator with extensive tuning.

- Trade-off:** Shower history reconstruction can be more complex.

### 💡 Why Angular Ordering?

Tracking shower history explicitly → Fits to new physics search

# Factorization Essentials

## Total Cross Section Formula

$$d\sigma_{i+j} \simeq \frac{\alpha_{\text{int}}(\tilde{q}^2)}{2\pi} \frac{d\tilde{q}^2}{\tilde{q}^2} dz P_{i\tilde{j} \rightarrow i+j}(z, \tilde{q}) \times d\sigma_{i\tilde{j}}$$

## Splitting Functions

- > Governs probability of parton emission (Altarelli-Parisi)
- > Resums collinear & soft emissions into parton showers
- > **Dependent on:** Spins, Masses, Ordering Variable

## Practical Impact

- ✓ Modular computation: Calculate pieces individually
- ✓ Validation: Isolate hard process from shower effects

## Factorization Theorem Workflow

1. PDF Convolution  
Parton Distribution Functions

×

2. Hard Scattering  
Matrix Element Calculation

×

3. Parton Shower/Hadronisation/Decay  
Resummation of QCD Effects

↓

**Final Hadronic Observables**

# BSM Showers in Herwig7

Model-Independent Framework for New Physics Simulation



## Phenomenology

Universal Splittings

- Only depends on spins, masses, and ordering variables of participating particles
- Based on the **Master Formula by Altarelli and Parisi** [NPB 126, 2 (1977)]

$$P_{0 \rightarrow 12}(z, \tilde{q}) = \frac{1}{2(q_0^2 - m_0^2)} \sum_{s_0, s_1, s_2} |\mathcal{M}_{s_0, s_1, s_2}|^2$$

- **Covers all spin combinations for emitters & spectators:**

spin-0

spin-1/2

spin-1

**Universal Coverage**



Hand Calculation



Coding



## In Herwig7

Automated & General

- **Full automation:** Reads BSM model information automatically from **UFO model files without manual intervention**
- Flexibility: can be interfaced with various matrix-element generators such as Herwig7's internal ME calculator and MG5, ...


**Automation is Key**

# Example Splitting: $q \rightarrow q \varphi$

Spin-Dependent Matrix Elements & General Coupling

Process: Fermion  $\rightarrow$  Fermion + Scalar

## Matrix Element Calculation



$$-i\mathcal{M} \left[ \begin{array}{c} \bar{u}(q_1) \\ \nearrow \\ \text{---} \\ \searrow \\ u(p) \end{array} \right] = \bar{u}(q_1) [-ig(\kappa + i\tilde{\kappa}\gamma_5)] u(p)$$

## Generalized Splitting Function

$$P_{f \rightarrow f' \phi}(z, \tilde{q}) =$$

$$\frac{g^2}{2} \left[ (\rho_+ |\kappa + i\tilde{\kappa}|^2 + \rho_- |\kappa - i\tilde{\kappa}|^2) \cdot [(1-z) - m_{2,t}^2] + (\rho_+ + \rho_-) [|\kappa|^2 (m_{0,t} + m_{1,t})^2 + |\tilde{\kappa}|^2 (m_{0,t} - m_{1,t})^2] + 2(\rho_+ - \rho_-) \Im(\kappa\tilde{\kappa}) [(1-2z)m_{0,t}^2 + m_{1,t}^2] \right],$$

[Lee, Masouminia, Seymour, Yang, JHEP 08 (2024) 064]



## Validation

### 1. SM Limit Recovery

$$g = g_W m_0 / 2m_W, \quad g_W = e / \sin \theta_W$$

$$m_0 = m_1, \quad \kappa = 1, \quad \tilde{\kappa} = 0, \quad \text{and} \quad \rho_0 + \rho_1 = 1,$$

$$P_{f \rightarrow f' \phi}^{SM}(z, \tilde{q}) = \frac{g_W^2}{8} \left( \frac{m_0}{m_W} \right)^2 \left[ (1-z) + \frac{4m_0^2 - m_2^2}{z(1-z)\tilde{q}^2} \right]$$

### 2. Performance test with FO


[JHEP 08 (2024) 064]

# Performance Test

Comparing Fixed-Order (FO) and Resummed Shower (RS) Approaches

## Fixed-Order (FO)


Full MadGraph5 Simulation



vs

## Resummed Shower (RS)

Herwig7 Parton Shower



### Expected Agreement

Comparing "HW( $Z'$ +SM PS) - HW(only  $Z'$  PS)" with "HW(only  $Z'$  PS) - MG( $Z'$ )" should show similar phenomena in the **collinear regime**.

# Z' shower in U(1)<sub>B-L</sub>

## Model Features

### Gauge Symmetry

$$U(1)_{B-L}$$

### Covariant Derivative

$$D_\mu = \partial_\mu + ig' Y_{B-L} Z'_\mu$$

### Lagrangian

$$\mathcal{L}_f = \sum_k i \bar{\psi}_k \gamma^\mu D_\mu \psi_k$$

$$\mathcal{L}_{\text{int}} \supset g'_1 Z'_\mu \sum_f \left( Y_{B-L}^f \bar{f} \gamma^\mu f \right)$$

## Properties

Vector coupling only

Anomaly Free

Mass = Gauge Eigenstate

No mixing

- The original splitting function

$$P_{f \rightarrow f'V}(z, \tilde{q}) = (|g_R|^2 \rho_+ + |g_L|^2 \rho_-) \left[ \frac{1+z^2}{1-z} (1 + m_{f,t}^2) - \frac{1+z}{1-z} m_{f',t}^2 - m_{V,t}^2 \right] \\ + (|g_R|^2 \rho_- + |g_L|^2 \rho_+) z m_{f,t}^2 - 2\text{Re}(g_L g_R^*) (\rho_+ + \rho_-) m_{f,t} m_{f',t},$$

- Symmetric property:  $g_L = g_R = g'_1(B - L)$  from the model

$$P_{f \rightarrow fV}(z, \tilde{q}) = \frac{g^2}{2} \left( \frac{1+z^2}{1-z} - 2m_{f,t}^2 - m_{V,t}^2 \right)$$


- The mass terms = dead-cone effect

# Performance test: $q \rightarrow q V$

Minimal extension of  $Z'$  from SM,  $U(1)_{B-L}$

Process: Fermion  $\rightarrow$  Fermion + Vector

## Basic process



@ 13.6 TeV  
w/  $m(Z') = 10$  GeV

## Validation results



**FO:**  $u\bar{u} \rightarrow d\bar{d}Z'$  (MG5)

**RS:**  $u\bar{u} \rightarrow d\bar{d}$  (MG5) +  $u \rightarrow uZ'$  (HW7)

Wonderful agreement @ collinear region

notable decrease @ non-collinear region

- ✓ No SM shower is included
- ✓ Only a single  $Z'$  emission is allowed in HW7



# BSM Parton Shower in Herwig7

Searching for New Physics Inside Jets with the Herwig 7  
Generalised Parton Shower

Joon-Bin Lee<sup>1</sup> • M.R. Masouminia<sup>3</sup> • Taehee Kim<sup>1</sup> • Un-ki Yang<sup>1</sup> • Michael H. Seymour<sup>2</sup>

<sup>1</sup> Seoul National University

<sup>2</sup> University of Manchester

<sup>3</sup> IPPP, Durham University



MCnet project



Marie Curie Actions



# Motivation



## Direction

First look

- Search  $Z' \rightarrow \mu\mu$  inside jets
- **Model-independent**, but pick up one **example model (B-L extension) for interpretation**  
 $SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$
- Signal samples are generated with HW BSM parton shower framework (previous work)
- **Search range:  $m(\mu\mu) = [5, 50]$  GeV**, i.e. above B hadron & below Z mass
- **Provide kinematic features of BSM PS & significance** under the current LHC setup

**Inclusive study is Key**

## Model Features

### Gauge Symmetry

$$U(1)_{B-L}$$

### Covariant Derivative

$$D_\mu = \partial_\mu + ig' Y_{B-L} Z'_\mu$$

### Lagrangian

$$\mathcal{L}_f = \sum_k i \bar{\psi}_k \gamma^\mu D_\mu \psi_k$$

$$\mathcal{L}_{\text{int}} \supset g'_1 Z'_\mu \sum_f \left( Y_{B-L}^f \bar{f} \gamma^\mu f \right)$$

### Properties

Vector coupling only

Anomaly Free

Mass = Gauge Eigenstate


No mixing

# Performance Test

Comparing Fixed-Order (FO) and Resummed Shower (RS) Approaches

## Fixed-Order (FO)

MadGraph5 Simulation



### Sample References


MG(Z')

MG(Z') + HW

VS

## Resummed Shower (RS)

Herwig7 Parton Shower



### Sample References

HW(only Z' PS)

HW(Z' + SM PS)

### ✓ Expected Agreement

“HW(Z'+SM PS)” – “MG(Z')+HW” / “HW(only Z' PS)” – “MG(Z')” should show similar phenomena @ collinear regime  
Comparing “HW(Z'+SM PS)” – “HW(only Z' PS)” shows SM PS effect


# Parton-level Validation Angular Separation $\Delta R(q, Z')$

## Previous Work

Only FSR-like diagrams considered in MG( $u\bar{u} \rightarrow d\bar{d} Z'$ ) sample  
**No SM Parton Shower** included, i.e. only  $Z'$  shower

## Current Work


Inclusive production of MG( $pp \rightarrow jj Z'$ ) events  
 Includes **full SM Shower +  $Z'$  shower** effects for **realistic** simulation



$m(Z') = 10$  GeV

Sample	MadGraph 5 (ME)		Herwig 7 (shower)	
	Hard process	$Z'$ shower	SM shower	
MG( $Z'$ )+HW	$pp \rightarrow Z'jj$	off	on	— (red solid)
MG( $Z'$ )	$pp \rightarrow Z'jj$	off	off	- - - (red dashed)
HW( $Z'$ +SM PS)	$pp \rightarrow jj$	on	on	— (black solid)
HW(only $Z'$ PS)	$pp \rightarrow jj$	first-emission only	off	- - - (black dashed)

$m(Z') = 5$  GeV



## Key Findings & Improvements

✔ **Good agreement** observed in collinear region ( $\Delta R < 1$ )

↑ HW( $Z'$ +SM PS) > HW(only  $Z'$  PS) as  $\Delta R \rightarrow 0$   
*Enhancement driven by cascade radiation effects*


# Parton-level Validation Energy Spectrum z (with $\Delta R < 1$ cut)

## Previous Work

Only FSR-like diagrams considered in MG( $pp \rightarrow Z' jj$ ) sample  
**No SM Parton Shower** included, i.e. only  $Z'$  shower

## Current Work


Inclusive production of MG( $pp \rightarrow Z' jj$ ) events  
 Includes **full SM Shower +  $Z'$  shower** effects for realistic simulation



$m(Z') = 10$  GeV

Sample	MadGraph 5 (ME)	Herwig 7 (shower)	
	Hard process	$Z'$ shower	SM shower
MG( $Z'$ )+HW	$pp \rightarrow Z' jj$	off	on
MG( $Z'$ )	$pp \rightarrow Z' jj$	off	off
HW( $Z'$ +SM PS)	$pp \rightarrow jj$	on	on
HW(only $Z'$ PS)	$pp \rightarrow jj$	first-emission only	off

$m(Z') = 5$  GeV



## Key Findings & Improvements

👍 Better agreement due to the  $\Delta R(q, Z') < 1$  cut

# Analysis Object-level Validation

Variable Transition

$\Delta R(q, Z')$

→

$\Delta R(j, \mu\mu)$


$z$  (Energy Fraction)

→

Muon Energy Fraction (MEF)


## Quark-Z' Separation: $\Delta R(q, Z')$

$\Delta R(q, Z')$  spectrum for  $m(Z') = 5$  GeV



## Jet-Dimuon Separation: $\Delta R(j, \mu\mu)$

$\Delta R(j, Z')$  spectrum for  $m(Z') = 5$  GeV



Sample	MadGraph 5 (ME)		Herwig 7 (shower)	
	Hard process	$Z'$ shower	SM shower	
MG(Z')+HW	$pp \rightarrow Z' jj$	off	on	— (solid red)
MG(Z')	$pp \rightarrow Z' jj$	off	off	- - - (dashed red)
HW(Z'+SM PS)	$pp \rightarrow jj$	on	on	— (solid black)
HW(only Z' PS)	$pp \rightarrow jj$	first-emission only	off	- - - (dashed black)

### Observation

More events observed as  $\Delta R(j, \mu\mu) \rightarrow 0$ , because anti-kt algorithm clusters jets around the collinear  $Z'$

# Analysis Object-level Validation

**Variable Transition**


z (Energy Fraction)




Muon Energy Fraction (MEF)

based on “MEF  $\approx 1 - z$ ” relation

## Energy fraction, z



## Muon Energy Fraction in a jet



**Observation**

MEF  $\rightarrow 0$ : Z' has large  $\Delta R$  with the quark; MEF  $\rightarrow 1$ : SM shower makes the final partons softer & non-collinear


# Kinematics of jets and muons Event Selection & Background Estimation

**Event selection:** OS dimuon pair with  $p_T > 52, 5 \text{ GeV}$  (for standrad trigger) &  $5, 5 \text{ GeV}$  (for scouting trigger) inside jets ( $\Delta R < 0.3$ )


**Background estimation:** Due to the large stat uncertainty, fit  $f(m_{\mu\mu}) = A m_{\mu\mu}^{-n} \times \frac{1}{2} \left[ 1 + \text{erf} \left( \frac{m_{\mu\mu} - \bar{m}}{\sqrt{2}\sigma} \right) \right]$

$\sim$  QCD cross section (power law)  $\leftarrow$   $p_T$  turn-on

Standard Trigger



Scouting trigger





Key Insight: Scouting trigger introduces no turn-on curve, improving limits @ low mass region

# Significance & Expected Limit

\* First-look utilizing Asimov significance without systematics consideration yet.

Method: Asimov (Stat. Only)



## Conclusion remarks



Can get  $3\sigma$  @  $m(Z') < 30 \text{ GeV}$  w/ Run-2+3 &  $5\sigma$  across the full search region with HL-LHC data



Because this study is the first look, we utilise the Asimov significance w/o consideration of systematics

# Systematics

Action item for the next work

from the paper

A detailed study of systematic uncertainties is beyond the scope of the present first-look analysis. Nevertheless, if the observable of interest is defined as a ratio of cross sections,

$$R(j \rightarrow Z') = \frac{\sigma(pp \rightarrow jj, j \rightarrow Z' \rightarrow \mu^+ \mu^-)}{\sigma(pp \rightarrow jj)},$$

the dominant uncertainties associated with QCD scale variations and parton distribution functions are expected to be significantly suppressed due to their substantial cancellation in the ratio. The remaining theoretical uncertainties are anticipated to originate primarily from EW Sudakov logarithms associated with  $Z'$  radiation and from the treatment of the bottom-quark mass scheme and fragmentation. These effects are expected to be moderate and, in principle, amenable to systematic control in more refined future studies.

# Summary & Outlook

## Achievements and Future Directions



### Achievements

Current Status

Completed

- ✓ First study using **Generalised BSM Shower** in Herwig7 and validate it against fixed-order calculations.
- ✓ Proposed novel search for  **$Z' \rightarrow \mu\mu$  inside jets** using non-isolated leptons.
- ✓ Demonstrated robust modeling of BSM radiation and **promising sensitivity** in 5–50 GeV range.
- ✓ Main feature of Herwig 7.4



### Future Work

Upcoming Tasks

Planned

- > **Systematics Study & Optimization of event selection**  
Include full systematic uncertainties and optimize event selection (b-tagging, vertexing).
- > **Model Interpretations**  
Extend analysis to other models: 2HDM+CS, hypercharge models, 3R, X models.
- > **Experimental Implementation**  
Collaborate with experimental groups (CMS/ATLAS) to implement search with real data.

# BSM Parton Shower in Herwig7

—  
END

Joon-Bin Lee<sup>1</sup> • M.R. Masouminia<sup>3</sup> • Taehee Kim<sup>1</sup> • Un-ki Yang<sup>1</sup> • Michael H. Seymour<sup>2</sup>

<sup>1</sup> Seoul National University

<sup>2</sup> University of Manchester

<sup>3</sup> IPPP, Durham University



MCnet project



Marie Curie Actions



# Event Generator = Bridge b/w Theory and Experiment

Connecting Experimental Reality with Theoretical Calculations



## Experiments

Real-world Data

- **Detector-level objects (tracks, clusters)**
- Complex environments: Pileup, electronic noise
- Trigger constraints & selection bias
- Complex final states with hadronization effects

**CMS / ATLAS / LHCb**



## Event Generators

~1,000 particles in  
a single event  
= **Ultra-high dimension  
integration**

**Analysis-Ready Events**



## Theory

Fundamental Physics


- **Lagrangians** & Feynman Rules
- Hard scattering Matrix Elements (ME)
- Parton Distribution Functions (PDFs)
- Resummation & perturbative calculations

**SM / BSM / EFT**

# Non-isolated lepton study

How to examine this collinear region?


Phenomenology → Experiment



# Non-isolated lepton study

How to examine this collinear region?

Phenomenology → Experiment



# Non-Isolated Leptons Inside Jets

Signal Characteristics & Physics Opportunity

New Physics Search



## Definition

Leptons contained physically within jet cones, characterized by **large isolation values** (sum of pT in cone > lepton pT). Distinct from standard isolated leptons used in SM precision measurements.



## Model Agnostic study

Can arise from the **parton shower process**, but this is just a single example w/o consideration of specific model. They are produced collinearly with other partons during the evolution of the jet.



## Motivation

1. Experimentally new observables
2. Complements existing searches @ low mass



## Physics Opportunity

Enables identification of  $Z' \rightarrow \mu\mu$  signals produced within jets. A previously unexplored phase space for light dark sector mediators.

# Signal Samples and Coupling Choice



Balancing Production Efficiency with Physics Accuracy



## Optimization Strategy


Coupling Trade-offs

- Goal:**  
 Maximize statistics while keeping multi-emission rate low.
- Too Large Coupling:**  
 Breaks down the  $\sigma \sim \alpha_{Z'}$  relation due to multiple emissions.
- Too Small Coupling:**  
 Inefficient production requires enormous computing resources.

$$N(1 Z') \propto \alpha_{Z'}$$

👉 **Final Choice:**  $N(>1 Z') / N(1 Z') \approx 0.01$

$$p_T(q) = 200\text{--}300 \text{ GeV}$$





**Figure 1.** Coupling-dependence study of  $Z'$  radiation in the parton shower for  $M(Z') = 5, 10, 20,$  and  $50 \text{ GeV}$ . Each panel shows the ratio of events containing more than one  $Z'$  boson to those containing exactly one  $Z'$  boson,  $N(>1 Z')/N(1 Z')$ , as a function of the fraction of di-jet events with exactly one  $Z'$  boson,  $N(1 Z')/N(\text{total})$  normalised by  $1/g_{qZ'}^2$ .

# Parton-level validation







- Good agreement @  $\Delta R(q, Z') < 1$
- $\text{HW}(Z'+\text{SM PS}) > \text{HW}(\text{only } Z' \text{ PS}) @ \Delta R \rightarrow 0$ 
  - ✓ enhanced by cascade radiation
- $\Delta R(q, Z') < 1$  cut is applied to  $z$  spectrum
  - ✓ Better agreement is achieved than the previous one

## Previous work (only FSR-like diagrams are considered)




# Analysis object-level validation

- Variables similar to those used in the parton-level validation are selected
- $\Delta R(q, Z') \rightarrow \Delta R(j, \mu\mu)$ 
  - ✓ Good agreement
  - ✓ More events @  $\Delta R(j, \mu\mu) \rightarrow 0$ , because jets are clustered by the anti-kt algorithm
- $z \rightarrow$  Muon Energy Fraction
  - ✓ Equivalent to  $1-z$  spectrum
  - ✓  $MEF \rightarrow 0$ :  $Z'$  has large  $\Delta R$  with the quark
  - ✓  $MEF \rightarrow 1$ : SM shower makes the final partons softer & non-collinear



# Analysis object-level validation



- $\Delta R(j, \text{leading/sub-leading } \mu)$
- Dip @  $\Delta R \rightarrow 0.4$  due to the jet reco algorithm
- In the later section, only muons with  $\Delta R < 0.3$  (conservative choice) are used to study those clustered as jets



**Figure 5.**  $\Delta R(\mu, \text{jet})$  distributions for  $M(Z') = 20$  GeV. The top (bottom) panels show the separation between the jet and the leading (subleading) muon. The left column covers the full range,  $\Delta R < 3.2$ , while the right column provides a zoomed view of the near-jet region,  $\Delta R < 0.5$ .

# Kinematics of jets and muons

## Jet-Muon Separation: $\Delta R(j, \mu)$



### ⚠ Reconstruction Artifact

- 📍 Dip at  $\Delta R \approx 0.4$  → boundary effects: **Jet Reconstruction Algorithm** (Anti-kt,  $R=0.4$ )
- 🔍 Muons near the jet boundary are less likely to be clustered or pass selection.

### ✔ Analysis Choice


To avoid this reconstruction artifact and ensure robust signal selection:

**Only muons with  $\Delta R < 0.3$  are used to study those clustered as jets.** (Conservative choice)

# Z' → μμ Inside Jets Search Strategy

Identifying Non-Isolated Leptons from BSM Showers

Methodology Overview



## Background Samples

QCD (pT-binned with PY8) and Top (ttbar + tW with POWHEG+PY8) are the dominant backgrounds.

QCD

Top

# The B-L Model Extension



## Simplest Gauge Extension

Anomaly-free model with conserved **B-L** (Baryon minus Lepton number).



## No Mixing with SM Bosons

Zero mixing in both mass & kinetic terms. No tree-level interaction between  $Z'$  and SM bosons.



## Pure Vector Coupling

The  $Z'$  boson is its own mass eigenstate with only vector couplings to fermions.



## Interpretation Role

Used as a benchmark for interpretation; our shower formalism remains **model-agnostic**.

### Lagrangian Features

#### Gauge Symmetry

$$U(1)_{B-L}$$

#### Covariant Derivative

$$D_\mu = \partial_\mu + ig' Y_{B-L} Z'_\mu$$

#### Properties

Anomaly Free

Mass = Gauge Eigenstate

# Significance & Expected Limit

\*First-look study utilizing Asimov significance without systematics consideration yet.

Method: Asimov (Stat. Only)

- Because this study is the first look, we utilise the Aimov significance w/o consideration of systematics
- We can get evidence-level significance @  $m(Z') < 30$  GeV with Run-2+3 data & discovery-level significance across the full search region with HL-LHC data



Current Era

**Run-2 + Run-3 Potential**



Future Era

**HL-LHC Discovery Reach**



Next Steps

**Refinement Plan**

Future iterations will include systematic uncertainties, optimized b-tagging, and improved dimuon vertex reconstruction.

## 📄 Conclusion remarks

\*First-look study utilizing Asimov significance without systematics consideration yet.



# Signal Samples and Coupling Choice

Balancing Production Efficiency with Physics Accuracy



## Optimization Strategy

Coupling Trade-offs

- **Goal:**

Maximize statistics while keeping multi-emission rate low.

$$N(1 Z') \propto \alpha_Z^2$$

- **Too Large Coupling:**

Breaks down the  $\sigma \sim \alpha_Z^2$  relation due to multiple emissions.

- **Too Small Coupling:**

Inefficient production requires enormous computing resources.

✓ **Final Choice:**  $N(>1 Z') / N(1 Z') \approx 0.01$



## Generated Samples

Simulation Datasets

- **HW(Z'+SM PS)**

Main signal sample with full BSM shower and SM shower effects.

- **HW(only Z' PS)**

Validation sample without SM shower (isolates BSM splitting).

- **MG(Z') + HW**

Matrix Element generation of Z' interfaced to Herwig.

- **HW(Z'+SM PS)**

Main signal sample with full BSM shower and SM shower effects.

Herwig 7.2

MadGraph5\_aMC

UFO Model

# Presentation Overview



## Fundamentals of Event Generators

How do we simulate proton-proton collision events?



## Parton Shower Algorithms

Understanding the role of parton shower & difference b/w PY8 and HW7.



## BSM extensions

Extending showers to BSM physics and why they are a crucial signal.




## What is the non-isolated lepton study?

Why should we study NIL? What we can observe?

# Event Simulation Workflow

From Proton-Proton Collision to Final-State Objects




## Factorization Theorem

Allows separating long-distance (low energy) physics from short-distance (high energy) interactions. Integrating all interactions at once is inefficient. Calculate each part individually.

# Event Simulation Workflow

From Proton-Proton Collision to Final-State Objects



## What Event Generators do



### Factorization Theorem

Allows separating long-distance (low energy) physics from short-distance (high energy) interactions. Integrating all interactions at once is inefficient. Calculate each part individually.