

TMD measurements in SIDIS with a jet at NLP

Max Jaarsma, Oscar del Rio, Ignazio Scimemi, Wouter Waalewijn
based on: [[arXiv:2507.03072](https://arxiv.org/abs/2507.03072)]

SCET 2026
Seoul





Cluster of Excellence
PRISMA++

Precision Physics,
Fundamental Interactions
and Structure of Matter





What to expect from this talk?



$$F_{PePp}^g(\phi_J, \phi_S) = \sum_i H_i \otimes F_i \otimes S_i \otimes J_i$$


- Motivation
- Introduction
- The SCET-II effective current at NLP
- Factorization of the hadronic tensor
- Soft-collinear overlap subtraction
- Final results + prediction
- Conclusion & outlook



Motivation

Motivation - TMD observables

- Probing the 3D structure of protons
- Sensitivity to a large range of scales
- Tests of universality of factorization
- Spin and azimuthal correlations
- Precision tests of the Standard Model




Adapted from seminar slides from J. Gaunt

Motivation - Power corrections in TMD observables

$$q_T \ll Q$$

- Cannot be ignored with improving experimental precision
- Many spin-dependent and azimuthal angle-dependent effects start at NLP
- Improve theoretical understanding of sub-leading power factorization



Ebert, Gao, Stewart (2021)

Vladimirov, Moos, Scimemi (2021)

Gamberg, Kang, Shao, Terry, Zhao (2022)

Talk by Johannes Michel at ESI workshop (2023)

Balitsky (2024)

Jets and TMD observables

The combination of TMD distributions with jet physics provides a powerful framework for probing hadronization and hadronic substructure.

- Jet-related factorization ingredients are perturbatively calculable for $q_T \gg \Lambda_{\text{QCD}}$
- This reduces the number of unknowns in a fit
- Symmetries reduce the number independent terms that appear in the cross section

Introduction to Power Corrections in TMD Observables

Introduction - SIDIS

■ $e^-(\ell) + h(P) \rightarrow e^-(\ell') + J(p) + X$

■ Process mediated by $\gamma^*(q)$


■ Measure the kinematic variables:

$$\mathbf{q}_T^2, \quad x = \frac{Q^2}{2q \cdot P}, \quad y = \frac{q \cdot P}{\ell \cdot P}$$

■ Include proton spin S

■ Measure the following angles:

$$\phi_J, \quad \phi_S$$



Introduction - SIDIS

■ $e^-(\ell) + h(P) \rightarrow e^-(\ell') + J(p) + X$

■ Process mediated by $\gamma^*(q)$


■ Measure the kinematic variables:

$$\mathbf{q}_T^2, \quad x = \frac{Q^2}{2q \cdot P}, \quad y = \frac{q \cdot P}{\ell \cdot P}$$

■ Include proton spin S

■ Measure the following angles:

$$\phi_J, \quad \phi_S$$



Introduction - What is the goal?

- Spin- and angular-dependence of cross section can be organized into form factors

$$\frac{d\sigma}{dx dy d\phi_J d\phi_S d\mathbf{q}^2} = \frac{\alpha_{\text{em}}^2 y}{8Q^2} \left\{ F_{UU} \frac{2 - 2y + y^2}{y^2} + F_{UU}^{\cos \phi_J} \cos \phi_J \frac{2(2 - y)\sqrt{1 - y}}{y^2} \right. \\ \left. + F_{LU}^{\sin \phi_J} \lambda_e \sin \phi_J \frac{2\sqrt{1 - y}}{y} + F_{UL}^{\sin \phi_J} S_{\parallel} \sin \phi_J \frac{2(2 - y)\sqrt{1 - y}}{y^2} + \dots \right\}$$

- Goal: Find factorized formulas for all form factors

$$F_{P_e P_p}^g(\phi_J, \phi_S) = \sum_i H_i \otimes F_i \otimes J_i \otimes S_i$$

Introduction - How do we get there?

- Introduce power counting

$$q_T \sim \lambda Q$$

- Identify relevant modes of momentum

$$\text{p-collinear} \sim (\lambda^2, 1, \lambda)$$


$$\text{J-collinear} \sim (1, \lambda^2, \lambda)$$

$$\text{soft} \sim (\lambda, \lambda, \lambda)$$

- Introduce background fields for modes

$$\phi = \phi + \phi + \phi + \phi$$

- Integrate out **off-shell modes**



Introduction - Recipe

- Start from the full-QCD hadronic tensor

$$W^{\mu\nu} = \int \frac{d^4b}{(2\pi)^4} e^{+iq\cdot b} \langle P | J^\mu(b) | J, X \rangle \langle J, X | J^\nu(b) | P \rangle$$

- Integrate out off-shell modes and insert the effective current

$$J^\mu \rightarrow J_{\text{eff}}^\mu = \sum_i C_i \otimes O_i \otimes S_i \otimes J_i$$

- Factorize the matrix elements and reshuffle color, spin and Lorentz indices

$$W^{\mu\nu} = \sum_i \Gamma_i^{\mu\nu} H_i \otimes F_i \otimes S_i \otimes J_i$$

The SCET-II Effective Current Operator at NLP

The effective current - What do we mean?

- Hard scattering governed by EM current


$$J^\mu = \bar{\psi} \gamma^\mu \psi$$

- Integrating out **off-shell** modes gives J_{eff}

$$J_{\text{eff}}^\mu = \int \mathcal{D}\psi \mathcal{D}\bar{\psi} \mathcal{D}A e^{iS_{\text{QCD}}[\phi]} e^{iS_{\text{int}}[\phi, \phi, \phi, \phi]} \\ \times [\bar{\psi} + \bar{\psi} + \bar{\psi} + \bar{\psi}] \gamma^\mu [\psi + \psi + \psi + \psi]$$

- Effective current can be written in the form

$$J_{\text{eff}}^\mu = \sum_i C_i \otimes \mathcal{O}_i \otimes \mathcal{O}_i \otimes \mathcal{O}_i$$



Standard approach

- Match QCD onto SCET-I current with

$$hc \sim (\lambda, 1, \lambda^{\frac{1}{2}})$$


$$s \sim (\lambda, \lambda, \lambda)$$

- Match **hard-collinear** operators onto **collinear** and **soft** operators.
- Matching is carried out by calculating matrix elements of the operators in partonic external states

Our approach

- Simultaneously integrate out all off-shell modes
- Happens on an operator level with position-space Feynman rules
- Effective operator is calculated directly, no matching involved

The effective current - Example of calculation



- Integrate out off-shell mode

$$J_{\text{eff}}^{\mu} \supset \bar{\psi} \gamma^{\mu} \int d^d w \frac{\Gamma(2 - \epsilon)}{2\pi^{d/2}} \frac{\psi}{(-w^2 + i0)^{2-\epsilon}} g A(w) \psi(w)$$

- Multipole expand

$$\Delta(w) A(w) \psi(w) \rightarrow \Delta(w) A(w^+ \bar{n}) \psi(w^- n) + \dots$$

- Apply Field equations to keep operator basis minimal

$$J_{\text{eff}}^{\mu} \supset \bar{\psi} \gamma^{\mu} \frac{1}{i\partial^-} g A^- \psi + n^{\mu} \bar{\psi} \left(A_T - \frac{i\partial_T}{i\partial^-} g A^- \right) \frac{1}{i\partial^+} \psi$$

The effective current - Match to SCET building blocks

- Match onto familiar SCET building blocks

$$\chi = W^\dagger \frac{\gamma^- \gamma^+}{2} \psi \qquad \mathcal{A}_T^\rho = W^\dagger [iD_T^\rho, W]$$


- This can be done order-by-order

$$\mathcal{A}_T^\rho = gA_T^\rho - \frac{i\partial_T^\rho}{i\partial^-} gA^- + \mathcal{O}(g^2)$$

- Two sets of building block operators for the soft fields

$$\begin{aligned} \Psi_n &= S_n^\dagger \psi & \mathcal{A}_{T,n}^\rho &= S_n^\dagger [iD_T^\rho, S_n] \\ \Psi_{\bar{n}} &= S_{\bar{n}}^\dagger \psi & \mathcal{A}_{T,\bar{n}}^\rho &= S_{\bar{n}}^\dagger [iD_T^\rho, S_{\bar{n}}] \end{aligned}$$

The effective current - Higher orders



- At higher orders, loops result in convolutions along the lightcone

$$C(\{x^+\}, \{y^+, y^-\}, \{z^-\}) \\ \times \mathcal{O}(x^+ \bar{n}) \mathcal{O}_n(y^- n) \mathcal{O}_{\bar{n}}(y^+ \bar{n}) \mathcal{O}(z^- n)$$

- Convolutions involving **Soft** operators can arise from hard-collinear loops.
- Wilson coefficients can be constrained from symmetries

The effective current - Summary of operators

- Usual LP operator
- Transverse derivative of LP building block
- Two building blocks operators
- $\bar{q}qg$ configuration

$$\bar{\chi}\gamma_T^\mu S_n^\dagger S_{\bar{n}}\chi$$

$$\bar{\chi}S_n^\dagger S_{\bar{n}}n^\mu \frac{i\partial_T}{i\partial^+}\chi$$

$$\bar{\chi}S_n^\dagger S_{\bar{n}} \frac{n^\mu}{i\partial^+} \mathcal{A}_T \chi$$

$$\bar{\chi}\gamma_T^\mu S_n^\dagger S_{\bar{n}} \mathcal{A}_T S_{\bar{n}}^\dagger S_n \frac{\gamma^-}{i\partial^-}\chi$$

The effective current - Summary of operators

- Soft gluon operator (constrained by $\partial_\mu J^\mu = 0$)
- Soft gluon operator (constrained by RPI)
- Soft gluon operator (unconstrained)
- Soft quark operator (unconstrained)

$$\bar{\chi} S_n^\dagger S_{\bar{n}} \mathcal{A}_{T,\bar{n}}^\rho \frac{n^\mu}{i\partial^+} \chi$$

$$\bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \frac{1}{i\partial^-} \mathcal{A}_{T,\bar{n}}^\rho \frac{i\partial_T^\rho}{i\partial^+} \chi$$

$$\bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \frac{i\partial_T}{i\partial^-} \mathcal{A}_{T,\bar{n}} \frac{i\partial_T^\rho}{i\partial^+} \chi$$

$$\bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \mathcal{A}_T \frac{\gamma^-}{i\partial^-} \Psi_{\bar{n}}$$

The effective current - Constraints from conservation

- Demand that the EM current is conserved up to power corrections

$$i\partial_\mu J_{\text{eff}}^\mu(x) = 0 + \mathcal{O}(\lambda^{\frac{7}{2}})$$

- Results in the following constraints

$$\begin{array}{lcl} n^\mu \bar{\chi} S_n^\dagger S_{\bar{n}} \frac{i\not{\partial}_T}{i\partial^+} \chi & \text{↻} & \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \chi \\ n^\mu \bar{\chi} S_n^\dagger S_{\bar{n}} \not{A}_{T,\bar{n}} \frac{1}{i\partial^+} \chi & \text{↻} & \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \chi \\ \frac{\bar{n}^\mu}{i\partial^-} \bar{\chi} \not{A}_T S_n^\dagger S_{\bar{n}} \chi & \text{↻} & \bar{\chi} \not{A}_T S_n^\dagger S_{\bar{n}} \frac{n^\mu}{i\partial^+} \chi \end{array}$$

The effective current - Constraints from RPI

- RPI: Choice of frame should not matter too much

$$\begin{array}{lll} \text{I:} & n \rightarrow n + \Delta_T & \bar{n} \rightarrow \bar{n} \\ \text{II:} & n \rightarrow n & \bar{n} \rightarrow \bar{n} + \bar{\Delta}_T \\ \text{III:} & n \rightarrow n e^\alpha & \bar{n} \rightarrow \bar{n} e^{-\alpha} \end{array}$$

- This connects the following operators

$$\begin{array}{ll} n^\mu \bar{\chi} S_n^\dagger S_{\bar{n}} \frac{i\not{\partial}_T}{i\partial^+} \chi & \text{↻} \quad \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \chi \\ \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \frac{1}{i\partial^-} \mathcal{A}_{T,\bar{n}}^\rho \frac{i\partial_T^\rho}{i\partial^+} \chi & \text{↻} \quad \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \chi \end{array}$$

The effective current - Summary

- Complete result for the SCET-II effective current at NLP
- Only 5 independent Wilson coefficients
- Constraints to be cross-checked

$$C_1 \otimes \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \chi$$

$$C_1 \otimes \bar{\chi} S_n^\dagger S_{\bar{n}} A_{\bar{n},T} \frac{n^\mu}{i\partial^+} \chi$$

$$C_1 \otimes \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \frac{1}{i\partial^-} A_{\bar{n},T}^\rho \frac{i\partial_T^\rho}{i\partial^+} \chi$$

$$C_2 \otimes \left(\frac{\bar{n}^\mu}{i\partial^-} - \frac{n^\mu}{i\partial^+} \right) \bar{\chi} A_T S_n^\dagger S_{\bar{n}} \chi$$

$$C_3 \otimes \bar{\chi} \gamma_T^\mu S_n^\dagger S_{\bar{n}} \frac{i\partial_T}{i\partial^-} A_{\bar{n},T} \frac{1}{i\partial^+} \chi$$

$$C_4 \otimes \bar{\chi} \gamma_T^\mu \gamma^- S_n^\dagger S_{\bar{n}} A_T \frac{1}{i\partial^-} \Psi_{\bar{n}}$$

$$C_5 \otimes \bar{\chi} \gamma_T^\mu \gamma^- S_n^\dagger S_{\bar{n}} A_T S_{\bar{n}}^\dagger S_n \frac{1}{i\partial^-} \chi_{\bar{n}}$$

Factorization of the Hadronic Tensor

Hadronic Tensor - Intermediate result

- The object of interest:

$$W^{\mu\nu} = \int \frac{d^4b}{(2\pi)^4} e^{+iq \cdot b} \langle P | J^\mu(b) | J, X \rangle \langle J, X | J^\nu(b) | P \rangle$$

- Direct insertion of the effective current results in large amount of terms
- Many terms vanish because of:
 - 1 Conservation of fermion number
 - 2 C, P and T symmetries
 - 3 Boost invariance of the vacuum

Hadronic Tensor - Intermediate result

- Boost invariance of the vacuum implies

$$0 = \langle 0 | \frac{1}{i\partial^-} \mathcal{A}_{T,\bar{n}}^\rho(b_T) S_{\bar{n}}^\dagger S_n(b_T) | X \rangle \langle X | S_n^\dagger S_{\bar{n}}(0) | 0 \rangle$$

$$0 = \langle 0 | \frac{i\partial_T^\sigma}{i\partial^-} \mathcal{A}_{T,\bar{n}}^\rho(b_T) S_{\bar{n}}^\dagger S_n(b_T) | X \rangle \langle X | S_n^\dagger S_{\bar{n}}(0) | 0 \rangle$$

$$0 = \langle 0 | S_n^\dagger S_{\bar{n}}(b_T) \left[\frac{1}{i\partial^-} \gamma^- \Psi_{\bar{n}}(b_T) \right]_\beta | X \rangle \langle X | \left[\frac{1}{i\partial^-} \bar{\Psi}_{\bar{n}}(0) \gamma^- \right]_\gamma S_{\bar{n}}^\dagger S_n(0) | 0 \rangle$$

Hadronic Tensor - Intermediate result

Leading-power

$$\langle \bar{\chi} | \chi \rangle \langle \chi | \bar{\chi} \rangle \langle S_n^\dagger S_{\bar{n}} | S_{\bar{n}}^\dagger S_n \rangle$$

Multipole correction

$$\langle \bar{\chi} | \chi \rangle \langle \chi | \bar{\chi} \rangle \langle i\partial^+ S_n^\dagger S_{\bar{n}} | S_{\bar{n}}^\dagger S_n \rangle$$

∂_T correction

$$\langle i\partial_T \bar{\chi} | \chi \rangle \langle \chi | \bar{\chi} \rangle \langle S_n^\dagger S_{\bar{n}} | S_{\bar{n}}^\dagger S_n \rangle$$

\mathcal{A}_T correction

$$\langle \bar{\chi} \mathcal{A}_T | \chi \rangle \langle \chi | \bar{\chi} \rangle \langle S_n^\dagger S_{\bar{n}} | S_{\bar{n}}^\dagger S_n \rangle$$

\mathcal{A}_T correction

$$\langle \bar{\chi} | \chi \rangle \langle \chi | \bar{\chi} \rangle \langle S_n^\dagger S_{\bar{n}} \mathcal{A}_{T,\bar{n}} | S_{\bar{n}}^\dagger S_n \rangle$$


Hadronic Tensor - Rapidity divergences

- Factorization ingredients contain rapidity divergences due to separation in rapidity
- Cancel between collinear, soft and anti-collinear
- Structure of cancellation more complicated at NLP

$$\langle P | \bar{\chi} \not{\partial}_T \chi | P \rangle$$

$$\langle P | \bar{\chi} \not{A}_T \chi | P \rangle$$


Ebert, Gao, Stewart (2021)
Vladimirov, Moos, Scimemi (2021)



Hadronic Tensor - Endpoint divergences

$$C_2 \otimes \bar{\chi} \mathcal{A}_T S_n^\dagger S_{\bar{n}} \chi \quad \Rightarrow \quad \int d\xi H_2(\xi, \dots) F(\xi, \dots) S(\dots) J(\dots)$$

↑
↑



$$F(\xi, \dots) \sim \frac{1}{\xi}$$

$$H_2(\xi, \dots) \sim \ln \xi$$

Liu, Neubert (2019)

Vladimirov, Moos, Scimemi (2021)

Talk by Johannes Michel at ESI workshop (2023)

Beneke, Garny, Jaskiewicz, Strohm, Szafron, Vernazza (2022)

Liu, Mecaj, Neubert, Wang (2022)

Hadronic Tensor - What to do with these divergences?


- Individual factorization ingredients contain rapidity divergences
- Convolutions between NLP ingredients result in endpoint divergences
- All divergences cancel in the cross section

Goal

Redefine ingredients such that all ingredients and convolutions are manifestly finite

Soft-Collinear Overlap (0-bin) Subtraction

Overlap subtraction - What's the deal?




- Soft and (anti-)collinear modes are separated by their virtualities
- Overlap region vanishes in dimensional regularization

Manohar, Stewart (2006)

Lee, Sterman (2006)

Idilbi, Mehen (2007)



- Soft and (anti-)collinear modes are separated by their rapidities
- Overlap region vanishes for many rapidity regulators, but not always

Overlap subtraction - Why do we need a proper formalism?

- At leading power, some rapidity regulators always result in vanishing overlap regions


Overlap subtraction

If the overlap always vanishes, why bother setting up a formalism?

- 1 Formulating factorization in a regulator-independent way
- 2 Overlap regions at sub-leading power do not necessarily vanish

Overlap subtraction - Method of regions treatment

- Method of regions: sum over all regions and subtract the overlaps
Beneke, Smirnov (1998)
- From a regions point of view, the convolution in ξ is part of the full loop integral
- The limit $\xi \rightarrow 0$ corresponds to the collinear gluon becoming soft



Overlap subtraction - The Field Theoretical Way

- 1 Split off a soft component from the collinear fields

$$\phi(x) \rightarrow \phi(x) + \phi(x)$$

- 2 Remove all leading-power soft-collinear overlap interactions by field redefinition

$$\psi(x) \rightarrow S_n(x)\psi(x) \quad A^\mu(x) \rightarrow S_n(x)A^\mu(x)S_n^\dagger(x)$$

- 3 Replace all collinear operators in the factorized formula by

$$\chi \rightarrow S_{\bar{n}}^\dagger S_n \chi + \Psi_{\bar{n}} \quad \mathcal{A}_T^\mu \rightarrow S_{\bar{n}}^\dagger S_n \mathcal{A}_T^\mu S_n^\dagger S_{\bar{n}} + \mathcal{A}_{T,\bar{n}}^\mu$$

- 4 Solve for the pure-collinear matrix element by inverting the transformation

Overlap subtraction - Example at leading-power

- 3 Apply overlap transformation to collinear operators

$$\langle P | \bar{\chi} | X \rangle \gamma^- \langle X | \chi | P \rangle \rightarrow \langle P | \bar{\chi} | X \rangle \gamma^- \langle X | \chi | P \rangle \frac{1}{N_c} \text{tr} [\langle 0 | S_n^\dagger S_{\bar{n}} | X \rangle \langle X | S_{\bar{n}}^\dagger S_n | 0 \rangle]$$

- 4 Invert transformation to obtain pure collinear transformation

$$\langle P | \bar{\chi} | X \rangle \gamma^- \langle X | \chi | P \rangle_{\text{pure}} = \frac{\langle P | \bar{\chi} | X \rangle \gamma^- \langle X | \chi | P \rangle}{\frac{1}{N_c} \text{tr} [\langle 0 | S_n^\dagger S_{\bar{n}} | X \rangle \langle X | S_{\bar{n}}^\dagger S_n | 0 \rangle]}$$

- This subtraction leads to the definition of physical TMD PDFs by

$$B \otimes B \otimes S \rightarrow \frac{B}{S} \otimes \frac{B}{S} \otimes S \equiv f \otimes f$$

Overlap subtraction - Example at next-to-leading-power

Overlap subtraction

At next-to-leading power also additive terms play a role

- Of particular importance for $\bar{\chi}\mathcal{A}\chi$ correlation functions

$$\begin{aligned}\langle P | \bar{\chi} \mathcal{A}(b_T) \chi(0) | P \rangle &\rightarrow \langle P | \bar{\chi} \mathcal{A}(b_T) \chi(0) | P \rangle \frac{1}{N_c} \text{tr} [\langle 0 | S_n^\dagger S_{\bar{n}}(b_T) S_{\bar{n}}^\dagger S_n(0) | 0 \rangle] \\ &+ \langle P | \bar{\chi}(b_T) \chi(0) | P \rangle \frac{1}{N_c} \text{tr} [\langle 0 | S_n^\dagger S_{\bar{n}}(b_T) \mathcal{A}(b_T) S_{\bar{n}}^\dagger S_n(0) | 0 \rangle]\end{aligned}$$

- Leads to subtraction term in definition of physical TMD PDFs

$$F_2 = \frac{F_2^{(0)}}{\sqrt{S}} - \frac{F_1 S_2}{\sqrt{S}}$$

Overlap subtraction - Effect on endpoint divergences

- Subtraction of the overlap should remove the leading $\xi \rightarrow 0$ behavior

$$F_2^{(0)}(\xi, \dots) \sim \frac{1}{\xi} + \mathcal{O}(\xi^0)$$

- At leading order the subtracted TMD PDF behaves as

$$F_2(\xi) = \frac{F_2^{(0)}}{\sqrt{S}} - \frac{F_1 S_2}{\sqrt{S}} \sim \mathcal{O}(\xi^0)$$

- Convolution in ξ is manifestly finite

$$\int d\xi H_2(\xi, \dots) F(\xi, \dots) = \text{finite}$$

The Hadronic Tensor at NLP

There are 3 types of contributions to the hadronic tensor at NLP

Leading-power

$$H_1 \otimes J_1 \otimes F_1$$

Kinematic correction

$$H_1 \otimes J_1 \otimes \partial_T^* F_1$$

Higher-twist correction

$$H_2 \otimes J_1 \otimes F_2$$

Final Results for the SIDIS Form Factors

Final results - Form factors

- Two form factors do not receive a correction at this order in q_T/Q ,

$$F_{UU,T} = f_1 \otimes J_1$$

$$F_{LL} = g_1 \otimes J_1$$

- Power corrections show up in angular modulations

$$F_{UU}^{\cos \phi_J} = -\frac{2|\mathbf{q}|}{Q} \left\{ \mathcal{J}_{0,0}[f_1 J_1] + \mathcal{J}_{1,1}[f_1 J_1'] \right. \\ \left. + \operatorname{Re}\left(\mathcal{J}_{1,1}^{(2)}[f_1 J_2]\right) + \operatorname{Im}\left(\mathcal{J}_{1,1}^{(2)}[f_2^\perp J_1]\right) \right\},$$

$$F_{LU}^{\sin \phi_J} = \frac{2|\mathbf{q}|}{Q} \left\{ \operatorname{Im}\left(\mathcal{J}_{1,1}^{(2)}[f_1 J_2]\right) - \operatorname{Re}\left(\mathcal{J}_{1,1}^{(2)}[f_2^\perp J_1]\right) \right\},$$

Final results - Phenomenological prediction

- Focus on one particular form factor, $F_{UL}^{\sin \phi_J}$, obtained by



$$d\Sigma|_{\sin \phi_J} \equiv \int_0^{2\pi} d\phi_J \frac{\sin \phi_J}{\pi} \int_0^{2\pi} d\phi_S \int_{0.01}^{0.95} dy \frac{d\sigma}{dx dy d\phi_J d\phi_S dq^2} \Big|_{\lambda_e, |S_\perp|=0}$$

- Jet functions can be calculated perturbatively
- Use a model (Gaussian in ξ) for the twist-3 TMD PDFs, we study 2:
 - ▶ Boer-Mulders model: $F_2 \sim h_1^\perp$
 - ▶ Sivers model: $F_2 \sim f_{1T}$
- Fits for these functions are available in the literature

Piloneta, Vladimirov (2024)

Bury, Prokudin, Vladimirov (2021)

Final results - Phenomenological prediction



- q_T dependence of the two models agree within error bands
- Large bands reflect the uncertainties in the non-perturbative TMD parameterization

Conclusions

Conclusions

- Constructed the SCET-II effective EM current to NLP
- Constructed the hadronic tensor for jet production in SIDIS
- Formulated 0-bin subtraction on an operator level
- After subtraction, only 3 types of contributions remained
 - ▶ Leading power F_1
 - ▶ Kinematic correction $\partial_T F_1$
 - ▶ Higher-twist correction F_2
- Derived factorization formulas for the form factors
- Phenomenological prediction for $F_{UL}^{\sin \phi_J}$

Thank you for your attention!

- Constructed the SCET-II effective EM current to NLP
- Constructed the hadronic tensor for jet production in SIDIS
- Formulated 0-bin subtraction on an operator level
- After subtraction, only 3 types of contributions remained
 - ▶ Leading power F_1
 - ▶ Kinematic correction $\partial_T F_1$
 - ▶ Higher-twist correction F_2
- Derived factorization formulas for the form factors
- Phenomenological prediction for $F_{UL}^{\sin \phi_J}$

Back-up

Introduction - What types of power corrections are there?

- Power corrections from kinematical variables

$$-g_T^{\mu\nu} L_{\mu\nu} = +2Q^2 \frac{2 - 2y + y^2}{y^2} - 4|\mathbf{q}|Q \cos(\phi_J) \frac{(2 - y)\sqrt{1 - y}}{y^2} + \mathcal{O}(|\mathbf{q}|^2)$$

- Power corrections from sub-leading derivative operators

$$\langle P | \bar{\chi}(b_T) \partial_T^{\rho} \chi(0) | P \rangle \Rightarrow \partial_T f_1(x, b_T)$$

- Power corrections from sub-leading field operators

$$\langle P | \bar{\chi}(b_T) \mathcal{A}_T \chi(0) | P \rangle \Rightarrow f_2(x, \xi, b_T)$$

Back-up slide: Position-space integration

$$\int dw^+ dw^- \frac{\mathcal{O}(w^+ \bar{n}) \mathcal{O}(w^- n)}{[-2w^+ w^- + i0]^\alpha} = \frac{-w^{1-\alpha} i\pi}{\Gamma(\alpha) \Gamma(1-\alpha)} \cdots \frac{1}{(i\partial^-)^\alpha} \mathcal{O} \frac{1}{(i\partial^+)^\alpha} \mathcal{O}$$

$$\partial_T^* = \partial_T + \frac{1}{2} \partial_T K(b_T) \ln\left(\frac{\zeta}{\bar{\zeta}}\right)$$

- Boer-Mulders model

$$g_{2L,\oplus/\ominus}^{\perp,q/\bar{q}}(x, \xi, \mathbf{b}^2) \simeq \frac{M^2 |\mathbf{b}|^2}{\sqrt{2} \pi^{3/2}} [\Theta(\xi - x_1) - \Theta(\xi - x_2)] e^{-\xi^2/2} h_1^\perp(x, \mathbf{b}^2)$$

- Sivers model

$$h_{1;f\leftarrow h}^\perp(x, \mathbf{b}^2) = A_f N_f \frac{(1-x)x^{\beta_q}(1+\epsilon_q x)}{n(\beta_q, \epsilon_q)} \exp\left(-\frac{r_0 + x r_1}{\sqrt{1 + r_2 x^2 \mathbf{b}^2}} \mathbf{b}^2\right)$$

Hadronic Tensor - Intermediate result

$$\begin{aligned}
 [\mathcal{W}^{\mu\nu}]_A^{(2,2)} &= (\gamma_T^\mu)_{\alpha\beta} (\gamma_T^\nu)_{\gamma\delta} \int \frac{d^2b}{(2\pi)^2} e^{+iq_T \cdot b_T} \\
 &\times \int dy^+ dy^- dz^+ dz^- e^{i(q^- z^+ + q^+ z^- - q^- y^+ - q^+ y^-)} C_1(y^+, y^-) C_1^*(z^+, z^-) \\
 &\times \int \frac{db^+}{2\pi} e^{+iq^- b^+} \langle P | \bar{\chi}_{i,\alpha}(b_T + b^+ \bar{n}) | X \rangle \langle X | \chi_{l,\delta}(0) | P \rangle \\
 &\times \int \frac{db^-}{2\pi} e^{+iq^+ b^-} \langle 0 | \chi_{j,\beta}(b_T + b^- n) | p, X \rangle \langle p, X | \bar{\chi}_{k,\gamma}(0) | 0 \rangle \\
 &\times \langle 0 | [S_n^\dagger S_{\bar{n}}(b_T)]_{ij} | X \rangle \langle X | [S_{\bar{n}}^\dagger S_n(0)]_{kl} | X \rangle .
 \end{aligned}$$

- E^n recombination scheme:

$$\mathbf{P}_{\text{Jet}}^{E^n} = \frac{k_1^0 + k_2^0}{(k_1^0)^n + (k_2^0)^n} \left[(k_1^0)^{n-1} \mathbf{k}_1 + (k_2^0)^{n-1} \mathbf{k}_2 \right] + \mathcal{O}\left(\frac{Q_T^2}{Q^2} Q_T\right)$$

- WTA recombination scheme (E^∞)

Bertolini, Chan, Thaler (2014)

$$\mathbf{P}_{\text{Jet}}^{\text{WTA}} = \Theta(k_1^0 - k_2^0) \frac{k_1^0 + k_2^0}{k_1^0} \mathbf{k}_1 + \Theta(k_2^0 - k_1^0) \frac{k_1^0 + k_2^0}{k_2^0} \mathbf{k}_2$$

- Energy-weighting enables soft-collinear factorization

— Refs

Overlap subtraction - Example at next-to-leading-power

$$\begin{aligned}
 & [\mathcal{F}_{q,21}^{\text{bare}}(x, \xi, b_T, \zeta)]_{\delta\alpha} \\
 &= i q^+ \int \frac{db_1^-}{2\pi} \frac{db_2^-}{2\pi} e^{-i\xi b_1^- q^+} e^{-i\xi b_2^- q^+} \\
 &\quad \times \left\{ \frac{\langle P | [\bar{\chi}(b_T + b_1^- n) \mathcal{A}_T(b_T + b_2^- n)]_{i,\alpha} | X \rangle \langle X | \chi_{i,\delta}(0) | P \rangle}{\sqrt{S(b_T, \zeta(\delta^+/q^+)^2)}} \right. \\
 &\quad \left. - \frac{\langle P | [\bar{\chi}(b_T + b_1^- n) \gamma_T^\rho]_{i,\alpha} | X \rangle \langle X | \chi_{i,\delta}(0) | P \rangle}{\sqrt{S(b_T, \zeta(\delta^+/q^+)^2)}} \right. \\
 &\quad \left. \times \frac{1}{N_c} \text{tr} \left[\langle 0 | S_n^\dagger S_n(b_T) \mathcal{A}_{T,n}^\rho(b_T + b_2^- n) | X \rangle \langle X | S_n^\dagger S_{\bar{n}}(0) | 0 \rangle \right] \right\}
 \end{aligned}$$

Overlap subtraction - Effect on rapidity divergences

- $i\partial_T \chi$ correction contains rapidity divergences not cancelled by \sqrt{S}

$$\begin{aligned} & \langle P | i\partial_T^\rho \bar{\chi}_\alpha(b_T + b^+ \bar{n}) | X \rangle \langle X | \chi_\delta(0) | P \rangle \rightarrow \\ & \langle P | i\partial_T^\rho \bar{\chi}_\alpha(b_T + b^+ \bar{n}) | X \rangle \langle X | \chi_\delta(0) | P \rangle \langle 0 | [S_n^\dagger S_{\bar{n}}(b_T)] | X \rangle \langle X | [S_{\bar{n}}^\dagger S_n(0)] | 0 \rangle \\ & + \langle P | \bar{\chi}_\alpha(b_T + b^+ \bar{n}) | X \rangle \langle X | \chi_\delta(0) | P \rangle \langle 0 | i\partial_T^\rho [S_n^\dagger S_{\bar{n}}(b_T)] | X \rangle \langle X | [S_{\bar{n}}^\dagger S_n(0)] | 0 \rangle . \end{aligned}$$

- Overlap subtraction introduces additional term involving $i\partial_T S_n$

$$\mathcal{A}_{T,n}^\rho S_n^\dagger S_{\bar{n}} = -\frac{1}{2} i\partial_T^\rho [S_n^\dagger S_{\bar{n}}] + \frac{1}{2} \left[S_n^\dagger S_{\bar{n}} \mathcal{A}_{T,\bar{n}}^\rho + \mathcal{A}_{T,n}^\rho S_n^\dagger S_{\bar{n}} \right] .$$

- Combining the \mathcal{A}_T correction, the above identity, and using C and P symmetry, one arrives at a manifestly finite expression

$$\partial_T B \otimes B \otimes S + \frac{1}{2} B \otimes B \otimes \partial_T S \rightarrow \partial_T^* \left(\frac{B}{\sqrt{S}} \right) \otimes \frac{B}{\sqrt{S}}$$